

Webinar Will Begin Momentarily

TODAY'S AGENDA:

- Welcome
- Speaker Introduction
- Presentation
- Q&A
- Closing

Orgain

Professional Education Series

Support. Inform. Educate. Empower.

Collagen Peptides in Clinical Practice: Mechanisms, Evidence, and Applications for Skin, Joint, and Bone Health



WEBINAR HOST:

Keith Hine, MS, RD

VP of Healthcare, Sports & Professional Education
Orgain, LLC



WEBINAR PRESENTER:

Shiloah Kviatkovsky, Ph.D, CISSN

Assistant Professor, Department of Orthopaedics
University of Arkansas for Medical Sciences

Funding from non-CPE revenue for CPE planning, development, review, and/or presentation has been provided by Orgain, LLC. The views expressed herein are those of the presenter and do not necessarily represent Orgain's views. The material herein is accurate as of the date it was presented and is for educational purposes only and is not intended to be medical advice. The material presented in this webinar, is not intended to be a substitute for professional medical advice, diagnosis, or treatment. You should seek the recommendation of a medical professional regarding a medical condition or treatment or before starting a new nutrition and/or health regimen. Reproduction or distribution of these materials is prohibited. ©2025 Orgain, LLC. All rights reserved.

Orgain, LLC is providing these webinars on an as-is basis and makes no representations or warranties of any kind with respect to the webinars. Orgain, LLC nor any of its directors, employees or other representatives will be liable for damages arising out of or in connection with the use of this document. This is a comprehensive limitation of liability that applies to all damages of any kind, including (without limitation) compensatory, direct, indirect or consequential damages, loss of data, income or profit, loss of or damage to property and claims of third parties.

Collagen Peptides in Clinical Practice: Mechanisms, Evidence, and Applications for Skin, Joint, and Bone Health

Evidence-Based Applications for Healthcare Professionals

Presenter:
Shiloah Kviatkovsky, PhD, MS, CISSN



Disclosures & Educational Framework

No Conflicts of Interest

This presentation is developed independently with no commercial sponsorship or financial relationships to disclose.

Grant/Research Support:

PB Leiner
Nestlé Health

• Consultant:


PB Leiner
Nestlé Health
Bonafide Health LLC

Evidence-Based Approach

Content is derived from peer-reviewed literature, systematic reviews, and randomized controlled trials published in scientific journals.

Clinical Application Focus

Information provided is intended for educational purposes to support evidence-based clinical decision-making in dietetic practice.

 **Disclaimer:** This presentation does not constitute medical advice. Clinical decisions should be individualized based on patient assessment, current evidence, and professional judgment.

Why collagen peptides are relevant to HCPs

High patient demand + widespread use

Patients routinely ask about collagen for skin aging, joint pain/OA, bone health, and recovery.

Evidence exists—but is heterogeneous

Data span RCTs and meta-analyses with differences in formulation, dose, duration, and endpoints.

Dietitians translate evidence to practice

Role includes expectation-setting, dosing/timing guidance, safety/allergen counseling, and product advice.

Best positioned as an adjunct, not a replacement

Collagen peptides complement total protein adequacy, resistance training/rehab, and condition-specific care plans.

Common questions you'll hear

“Does collagen actually work?”

“How much should I take—and for how long?”

“Is one type better (type I vs type II)?”

“What's the difference between collagen and whey protein?”





Learning Objectives

01

Differentiate collagen structures and bioavailability

Distinguish between collagen types, native collagen, gelatin, and hydrolyzed collagen peptides, including their absorption characteristics and clinical implications.

02

Discuss potential mechanisms of action

Describe potential mechanisms of action by which collagen peptides may improve skin, joint, and bone health.

03

Evaluate clinical evidence across outcomes

Analyze the strength of evidence for collagen peptides in skin health, joint function, bone metabolism, and emerging applications in clinical populations.

04

Apply practical dosing and timing strategies

Implement evidence-based protocols for collagen peptide supplementation, including optimal dosing, timing relative to exercise, and combination with cofactors like vitamin C.

05

Counsel patients with realistic expectations

Apply practical, evidence-based recommendations when counseling patients about collagen.

Agenda

01

Collagen 101 + Mechanisms of Action

Exploring collagen in the body, in supplement form, and potential pathways: likely more than one mechanism at work

03

Joint Health Data

Clinical evidence for collagen peptides in joint function, pain management in OA, and tendon and ligament health

05

Additional Clinical Applications

Emerging research findings on collagen peptides and muscle, wound healing, and perioperative care

02

Collagen and Skin

Understanding what's really happening at the cellular and tissue level

04

Bone Health Data

Research findings on collagen peptides and bone density, turnover, and strength

06

Summary and Ending Remarks

Overview of dosing strategies and clinical applications for HCPs






SECTION OVERVIEW

Collagen 101 + Mechanisms of Action

What is collagen and how does it work?

Understanding Collagen: The Body's Structural Protein



1/3

Total Body Protein

Collagen comprises one-third of the body's total protein content

70%

Skin Composition

The majority of skin's dry weight

80%

Tendons & Ligaments

Provides tensile strength and flexibility

30%

Bone Structure


Forms the organic matrix of bone tissue

Collagen is the structural building block of our body

Collagen is also essential in eyes, ear and nose cartilage, teeth, hair, lungs, blood vessels, muscles, and the gastrointestinal tract.

Clinically Relevant Collagen Types

Over **28 collagen** types exist in the human body, but three dominate clinical applications based on tissue distribution and functional significance.



1

Type I Collagen

Location: Skin, bone, tendons, ligaments, cornea

Function: Provides tensile strength and structure; most abundant (90% of body collagen)

Clinical focus: Skin elasticity, bone density, wound healing

2

Type II Collagen

Location: Articular cartilage, intervertebral discs, vitreous humor

Function: Provides compressive resistance in cartilage

Clinical focus: Joint health, osteoarthritis management

3


Type III Collagen

Location: Skin, blood vessels, internal organs, reticular fibers

Function: Supports tissue architecture alongside Type I






Clinical focus: Skin structure, cardiovascular tissue integrity



Collagen Forms: Bioavailability & Clinical Application



📌 **Clinical Pearl:** Hydrolyzed collagen peptides demonstrate superior absorption compared to gelatin or native collagen. Plasma concentrations of hydroxyproline-containing peptides peak 1-2 hours post-ingestion and remain elevated for up to 4 hours, supporting tissue accumulation.

Hydrolyzed Collagen Peptides

TYPE OF COLLAGEN SUPPLEMENT	
	Bovine Collagen
	Pig Collagen
	Chicken Collagen
	Fish and Marine Collagen
	Egg Shell Membrane Collagen



Absorption and Bioavailability: What the Evidence Shows

FACT

WE DON'T BREAK DOWN ALL PROTEIN INTO SINGLE AMINO ACIDS

- Like any protein, collagen can be digested into single amino acids, or into small peptide chains
- Collagen di- and tri-peptides often absorbed whole
- Even slightly longer chains of 20-50 amino acids (2k-5kDa) can remain intact ... and be absorbed






Image Credit: Absolute Collagen

Collagen Stewardship Alliance
fueled by
INGREDIENT
TRANSPARENCY
CENTER




Mechanisms of Action: 3 Hypotheses

1. A SUPPLY OF AMINO ACIDS



2. BIOLOGIC MESSENGERS



Collagen peptides stimulate activity of cells that produce new collagen and degrade old collagen fibrils:

Fibroblasts, Tenocytes, Chondrocytes and Osteoblasts

3. INDUCERS OF ORAL TOLERANCE

Oral Tolerance: Reduced immune response to previous antigens.

Collagen breakdown products in joints may serve as antigens.

Repeated ingestion of collagen peptides may "tolerize" the body to collagen, thus reducing the immune response in the joints.

*for undenatured type II collagen only


CLINICAL EVIDENCE

Evidence — Skin



Hallmarks of skin damage

Loss of physiological function and structural integrity



Collagen Peptides and Skin RCT

In a trial of women ages 45-65y
(n=108)

- 2.5 g of collagen peptides vs. placebo
- 8 weeks
- Compared to the control group:
 - CP significantly reduced wrinkles ($p < .05$)
 - CP increased synthesis of skin procollagen type I by 65% and elastin by 18%





Fig. 1. a Orally administered BCP led to a statistically significant reduction of eye wrinkle volume after 4 and 8 weeks of treatment. Over the same period of time, eye wrinkle volume in the placebo-treated group increased continuously (mean \pm SEM; $n = 57$; * $p < 0.05$, ** $p < 0.01$). **b** Visible reduction of eye wrinkle volume after 8 weeks of BCP intake. Exemplary pictures of 2 participants of the active agent group before (left) and after (right) treatment.


What the Systematic Reviews Show



Meta-Analysis Conclusions

- **Hydration:** Significant improvement across multiple RCTs (SMD 0.4–0.8)
- **Elasticity:** Moderate but consistent gains, especially in women over 35
- **Wrinkles:** Visible reduction in eye area; effects stronger with longer supplementation
- **Safety:** Excellent tolerability profile; adverse events rare and mild
- **Funding bias:** Industry-funded trials show larger effects, but independent studies still demonstrate benefit

Collagen in Photoaging



Skin Health Outcomes: Clinical Evidence Summary



Primary Outcomes Demonstrated

- **Skin elasticity:** Significant improvements in elasticity measures (R2, R5, R7) after 8-12 weeks
- **Hydration:** Increased skin hydration and reduced transepidermal water loss
- **Wrinkle reduction:** Decreased wrinkle depth and improved skin surface structure
- **Dermal density:** Ultrasound and biopsy studies show increased dermal collagen density

2.5-15

Dose Range (grams/day)

Most studies use 2.5-10 g daily; lower doses effective for skin outcomes

8-12

Weeks to Visible Effects

Clinically meaningful changes typically observed by 8-12 weeks of consistent use

12


Studies in Meta-Analysis

Systematic reviews include 12+ RCTs showing consistent benefits across populations

Who Benefits Most: Women ages 35-65, those with photoaged skin, individuals with reduced dietary protein or collagen intake. Effects more pronounced in older adults with baseline collagen decline.

CLINICAL EVIDENCE

Evidence — Joints/OA




Joint Physiology

TWO CONTRIBUTORS TO JOINT PAIN:

Joints lose collagen and other matrix components that maintain structure and provide cushion between bones, joints and ligaments.

Joints produce inflammatory mediators, that lead to swelling
stiffness and discomfort.



Recent RCT Spotlight: Low-Molecular-Weight Collagen



Park et al. 2025 — Key Findings

Design: 6-month RCT in knee OA patients (n=80) 40-75 yrs.

Intervention: 3 g/day low-molecular-weight collagen peptides vs. placebo

Primary outcome WOMAC pain and physical function subscales, as well as WOMAC total improved significantly. No serious adverse events.

Clinical takeaway: Low-MW peptides may enhance absorption and efficacy. Results align with broader meta-analysis trends.

Collagen Peptides vs. UC-II: Clinical Positioning

Collagen Peptides (Hydrolyzed)

- **Dose:** 5–10 g/day
- **Mechanism:** Provides amino acids (glycine, proline, hydroxyproline) for cartilage synthesis
- **Evidence:** Multiple RCTs and meta-analyses; moderate-quality evidence
- **Best for:** General joint health, athletes, mild-moderate OA



UC-II (Undenatured Type II Collagen)

- **Dose:** 40 mg/day
- **Mechanism:** Oral tolerance; modulates immune response in joint tissue
- **Evidence:** Smaller number of RCTs; promising but less meta-analysis data
- **Best for:** Moderate-severe OA, patients seeking lower-dose option

Both are safe and evidence-based. Choice depends on patient preference, cost, and OA severity. Not typically combined.

Meta-Analysis Summary: Clinical Interpretation

Pain Reduction

Pooled analysis shows significant VAS score improvement (mean difference: -10 to -15 mm). Effect size comparable to low-dose NSAIDs in mild-moderate OA.

Functional Improvement

WOMAC physical function scores improve moderately. Patients report easier stair climbing, walking, and daily activities.

Safety Profile

Adverse events minimal. GI symptoms rare. No drug interactions reported. Well-tolerated long-term.

Evidence strongest for knee OA. Hand and hip OA data emerging but less robust. Heterogeneity exists across trials in dose and peptide type.



Joint Health: Osteoarthritis Pain & Function

Evidence Base for Collagen Peptides in OA

Multiple randomized controlled trials and systematic reviews support the use of collagen peptides for osteoarthritis symptom management. Evidence is strongest for knee OA, with emerging data for hip and hand joints.

1

Pain Reduction

Significant reduction in VAS and WOMAC pain scores compared to placebo. Effect size modest but clinically meaningful, particularly for mild-moderate OA.

Recommended Dosing

Standard dose: 10 g/day hydrolyzed collagen

Alternative: 40 mg/day undenatured type II collagen (UC-II) for immune-mediated mechanism

Duration: Minimum 12 weeks; optimal benefits at 24+ weeks

2

Functional Improvement

Improvements in WOMAC function subscale, stair climbing ability, and joint mobility. Effects emerge gradually over 12-24 weeks.


Clinical Positioning


Adjunctive therapy for OA management. Consider for patients seeking non-pharmacologic options, those with GI intolerance to NSAIDs, or as part of comprehensive joint health strategy.

3

Cartilage Biomarkers

Reduced CTX-II (cartilage degradation marker) and increased type II collagen synthesis markers suggest protective effects on cartilage metabolism.






Joints — Tendon & Ligament

CONNECTIVE TISSUE SUPPORT

Key Evidence: Collagen Synthesis Boost



Shaw et al. 2017 Study

Landmark research demonstrated that 15 g gelatin (collagen source) plus vitamin C, taken 1 hour before 6 minutes of intermittent rope-skipping exercise, doubled collagen synthesis markers in serum.

Mechanism: Vitamin C is essential for hydroxylation of proline and lysine residues in collagen. Exercise provides mechanical signal. Together, they optimize collagen fibrillogenesis.

Clinical Implication: Strategic timing matters. This synergy model informs rehabilitation protocols and injury prevention strategies in athletic populations.

Loading Synergy: The Pre-Exercise Model

01

Collagen Timing

Consume 10–15 g collagen peptides or gelatin consumed **30–60 minutes before exercise** or rehabilitation activity (Shaw et al., 2017; Lis et al., 2019)

03


Vitamin C Co-ingestion

50 mg vitamin C consumed with collagen. Essential cofactor for hydroxylation of proline residues during collagen synthesis (Shaw et al., 2017)


02

Mechanical Loading

Exercise or rehab session provides mechanical stimulus to target tissues, enhancing collagen synthesis when combined with pre-exercise intake

 **Emerging Strategy:** This tripartite approach—pre-exercise collagen + vitamin C + loading—maximizes collagen synthesis in target tissues.





Evidence — Bone Health

CLINICAL APPLICATIONS

BONE COMPOSITION

MINERALS EMBEDDED IN A COLLAGEN MATRIX

50-70 % Mineral,
hydroxyapatite
crystal

20-40 % Organic
Matrix (90% Type I
collagen)

5-10 % Water

Small amount of
lipid


Bones and the Remodeling Process

OSTEOBLASTS are involved in bone formation and mineralization

OSTEOCLASTS are involved in bone breakdown or resorption.

BONE MINERALIZATION is the process where minerals are added to the organic matrix of bone.

BONE REMODELING (or bone metabolism) is a continuous process where mature bone tissue is removed from the skeleton (also called bone resorption) and new bone tissue is formed.



Key RCT: Postmenopausal Women



Dose

5 grams daily of specific collagen peptides (CP) for 12 months



BMD Outcomes

Significant increase in spine BMD (+2.3%) vs. placebo; favorable trend in femoral neck



Marker Changes

P1NP increased (formation), CTX decreased (resorption) — net anabolic signal



Clinical Pearl: The König 2018 study used a specific bioactive collagen peptide formulation. Response may vary with peptide source and molecular weight profile.



Meta-Analysis: Clinical Positioning

What the Evidence Shows

Recent meta-analysis (Sun et al. 2025) pooled multiple RCTs and found:

- Consistent improvements in bone formation markers across studies
- Modest but significant BMD gains in spine and hip regions
- Best results with 5–10g daily for ≥ 12 months
- Effects more pronounced in postmenopausal populations

Current Limitations


Heterogeneity in peptide types, study duration variability, and limited data in younger populations. Not a replacement for established osteoporosis therapies.

Effective Dose Range

5-10g

Months Needed

12+





Bone Outcomes: Where Collagen Fits

Bone Mineral Density (BMD)

Collagen peptides may support BMD maintenance in postmenopausal women and older adults through stimulation of osteoblast activity and modulation of bone remodeling pathways.

Evidence suggests modest improvements in spine and femoral neck BMD with consistent supplementation over 12+ months.

Bone Turnover Markers

More consistent effects observed on biochemical markers including:

- P1NP (bone formation marker) — typically increases
- CTX (bone resorption marker) — often decreases
- Osteocalcin and other regulatory proteins

Marker changes suggest favorable shift in bone metabolism balance.

Evidence — Muscle & Sarcopenia

EMERGING APPLICATION



Collagen + Resistance Training

What It May Improve

- Fat-free mass gains in older adults (sarcopenic populations)
- Functional strength outcomes when combined with structured resistance training
- Body composition shifts (modest fat mass reduction)
- Connective tissue within and around muscle

What It Doesn't Replace

- High-quality, leucine-rich protein sources (whey, casein, etc.)
- Complete amino acid profile for muscle protein synthesis
- Gold-standard post-exercise recovery nutrition

📌 **Low Leucine Caveat:** Collagen is naturally low in leucine (~2–3% vs. 10% in whey), limiting its direct mTOR activation. Best used as an adjunct, not a replacement for complete proteins.



CLINICAL APPLICATIONS

Wound Healing & Pressure Injury Management

Evidence-based strategies for integrating collagen peptides into nutrition care plans for wound healing and tissue repair.



RCT Evidence: Collagen Hydrolysate for Pressure Ulcer Healing



Lee SK et al., 2006

Adv Skin Wound Care – 89 nursing home residents with stage II–IV pressure ulcers. Collagen hydrolysate group showed improved healing vs. placebo at 8 weeks.



Yamanaka H et al., 2017

J Nutr Intermed Metab – Improved DESIGN-R scores and faster healing in elderly patients receiving collagen peptide supplementation alongside standard care.



Sugihara F et al., 2018

Sci Rep – Double-blind placebo-controlled trial demonstrated enhanced pressure ulcer healing rate and reduced wound size with collagen peptide intervention.

📌 Note: All trials provided collagen peptides **in addition to** adequate baseline nutrition—not as a replacement for protein or calories.

PERIOPERATIVE & ORTHOPEDIC

Perioperative & Orthopedic Recovery

Emerging evidence for collagen peptides as a supportive adjunct in surgical recovery and tissue repair.





Emerging Evidence: Supportive Adjunct Role

ACL Rehabilitation Support

López-Vidriero E et al. (2019) found collagen peptide supplementation combined with progressive rehabilitation improved functional outcomes and patient-reported knee function post-ACL reconstruction.

Framing as Adjunctive

Current evidence suggests collagen peptides may support tendon, ligament, and joint cartilage remodeling—but always **alongside** physical therapy, adequate protein intake, and standard postoperative care.

What We Don't Know Yet


Large-scale RCTs in diverse surgical populations are lacking. Effect sizes are modest. Mechanism (direct vs. indirect via amino acid profile) remains under investigation.

📌 Caution: Avoid marketing language. Frame collagen peptides as **potentially supportive**, not curative or essential.

MECHANISMS, EVIDENCE, & PRACTICE

Summary & Clinical Takeaways

Turning evidence into practical dosing, timing, and counseling guidance for HCPs




Summary of Mechanisms of Action

Substrate Provision
Supplies amino acids (glycine, proline, hydroxyproline) as building blocks for endogenous collagen synthesis

Matrix Protection
May inhibit matrix metalloproteinases (MMPs) that break down collagen, preserving tissue integrity

Vitamin C Synergy

Vitamin C is essential for hydroxylation of proline and lysine residues, stabilizing collagen structure. Co-supplementation (50-100 mg) enhances collagen synthesis.



Cell Signaling

Bioactive peptides stimulate fibroblasts and chondrocytes to increase ECM production via receptor-mediated pathways

Mechanical Loading

Synergy with exercise: mechanical stress + peptide availability enhances tissue remodeling response

Exercise Timing

Consuming collagen peptides 30-60 minutes before mechanical loading optimizes tissue availability during the anabolic window of tissue repair.

Clinical Implementation: Dosing, Timing & Counseling Applications



Evidence-Based Dosing by Indication

Skin: 2.5-10 g/day | **Joints:** 10 g/day or 40 mg UC-II | **Bone:** 5-10 g/day | **Tendon/ligament:** 15 g/day pre-exercise



Optimal Timing Strategies

General use: any consistent time. For musculoskeletal rehab: 30-60 min before loading/exercise. With vitamin C (50-100 mg) to enhance synthesis.




Setting Realistic Expectations

Educate on timeline: 4-8 weeks for **skin**, 12-24 weeks for **joint** benefits. Emphasize consistency. Not a replacement for comprehensive nutrition or medical therapy.



Product Quality Considerations

Recommended products
Verify collagen source and hydrolysis method and type. Check for added sugars or unnecessary ingredients.

 **Safety Profile:** Collagen peptides demonstrate excellent safety and tolerability. Mild GI effects (fullness, mild nausea) reported rarely. No significant drug interactions identified. Consider individual allergies to source material (bovine, marine, chicken).

Dosing and Timing Summary

Dose ladder

2.5 g













5 g

10 g

15 g

15 g

Most-studied daily dose range
*goal dependent

Goal	Dose (daily)	Trial Duration	Best paired with	What to tell patients
Skin	2.5–10 g	 8–12 weeks	 Adequate protein + skincare	 30–60 min pre-session most commonly used  8–12 weeks
OA pain/function	5–15 g	 8–24 weeks	 Strengthening/PT + weight management	
Bone health	~5 g	 6–12 months	 Resistance training + Ca/Vit D	Supportive adjunct; not a substitute for osteoporosis care
Tendon/ligament rehab	10–15 g pre-loading*	 weeks–months	 Loading/rehab + Vit C	Most compelling when timed around rehab/loading.
Wound healing/pressure injury	variable**	 4–12 weeks	 Energy + protein sufficiency + standard wound care	Consider adjunct in high-risk patients; avoid overpromising.

Key Takeaways & Questions

Evidence-Based, Not Hype-Based

Collagen peptides show promise for wound healing and tissue recovery—but always as part of comprehensive nutrition and rehab care.


Individualize & Monitor

Assess baseline nutrition, set realistic expectations, and use validated tools to track outcomes. Not every patient will benefit equally.

Quality Matters

Trusted products, screen for allergens, and counsel on appropriate dosing and timing for target tissues.

Questions?



Thank you for joining us today.

*RDNs and NDTRs, please complete the
short survey after the webinar to
download a CPEU certificate!*

Orgain.

Professional Education Series

Support. Inform. Educate. Empower.



WEBINAR HOST:

Keith Hine, MS, RD

VP of Healthcare, Sports & Professional Education
Orgain, LLC

keith.hine@orgain.com



WEBINAR PRESENTER:

Shiloah Kviatkovsky, Ph.D, CISSN

Assistant Professor, Department of Orthopaedics
University of Arkansas for Medical Sciences

skviatkovsky@uams.edu



ORGAIN INQUIRIES:

medinfo@orgain.com

Funding from non-CPE revenue for CPE planning, development, review, and/or presentation has been provided by Orgain, LLC. The views expressed herein are those of the presenter and do not necessarily represent Orgain's views. The material herein is accurate as of the date it was presented and is for educational purposes only and is not intended to be medical advice. The material presented in this webinar, is not intended to be a substitute for professional medical advice, diagnosis, or treatment. You should seek the recommendation of a medical professional regarding a medical condition or treatment or before starting a new nutrition and/or health regimen. Reproduction or distribution of these materials is prohibited. ©2025 Orgain, LLC. All rights reserved.

Orgain, LLC is providing these webinars on an as-is basis and makes no representations or warranties of any kind with respect to the webinars. Orgain, LLC nor any of its directors, employees or other representatives will be liable for damages arising out of or in connection with the use of this document. This is a comprehensive limitation of liability that applies to all damages of any kind, including (without limitation) compensatory, direct, indirect or consequential damages, loss of data, income or profit, loss of or damage to property and claims of third parties.